

Rehabilitation Protocol: Arthroscopic Anterior Shoulder Stabilization (Bankart Repair)

Name:	2.1.10	
Date:	Date of Surgery:	
Diagnosis:		_

Phase I (Weeks 0-4)

- Sling immobilization at all times except for showering and rehab under guidance of PT
- Range of Motion PROM → AAROM → AROM as tolerated
 - Restrict motion to 90° of Forward Flexion, 45° of Abduction, 30° of External Rotation and Internal Rotation to stomach
- Therapeutic Exercise
 - Elbow/Wrist/Hand Range of Motion
 - o Grip Strengthening
 - Isometric Abduction, Internal/External Rotation exercises with elbow at side
- · Heat/Ice before and after PT sessions

Phase II (Weeks 4-6)

- Discontinue sling immobilization
- Range of Motion Increase Forward Flexion, Internal/External Rotation to full motion as tolerated
- Therapeutic Exercise
 - Advance isometrics from Phase I to use of a theraband within AROM limitations
 - Continue with Elbow/Wrist/Hand Range of Motion and Grip Strengthening
 - Begin Prone Extensions and Scapular Stabilizing Exercises (traps/rhomboids/levator scapula)
 - o Gentle joint mobilization
 - Modalities per PT discretion

Phase III (Weeks 6-12)

- Range of Motion Progress to full AROM without discomfort
- Therapeutic Exercise Advance theraband exercises to light weights (1-5 lbs)
 - 8-12 repetitions/2-3 sets for Rotator Cuff, Deltoid and Scapular Stabilizers
 - Continue and progress with Phase II exercises
 - Begin UE ergometer
 - Modalities per PT discretion

Phase IV (Months 3-6)

- Range of Motion Full without discomfort
- Therapeutic Exercise Advance exercises in Phase III (strengthening 3x per week)

- o Sport/Work specific rehabilitation
- Return to throwing at 4.5 months
 Return to sports at 6 months if approved
 Modalities per PT discretion

Comments:				
Frequency: Signature:	_ times per week	Duration:	weeks Date:	