

Rehabilitation Protocol: Meniscus Allograft Transplantation

Name: Date:	Date of Surgery:	
Diagnosis:	- 1	

Phase I (Weeks 0-8)

- Weightbearing:
 - Weeks 0-2: Partial Weightbearing (up to 50%)
 - Weeks 2-6: Advance to WBAT with crutches (d/c crutches at 4 weeks post-op if gait normalized)
- Hinged Knee Brace: worn for 6 weeks post-op
 - Locked in full extension for ambulation and sleeping remove for hygiene
 (Week 1)
 - Locked in full extension for ambulation
 – remove for hygiene and sleeping
 (Week 2)
 - Set to range from 0-90° for ambulation- remove for hygiene and sleeping (Weeks 3-6)
 - Discontinue brace at 6 weeks post-op
- Range of Motion PROM → AAROM → AROM as tolerated
 - Weeks 0-2: Non-weightbearing 0-90°
 - Weeks 2-8: Full non-weightbearing ROM as tolerated progress to flexion angles greater than 90°
- Therapeutic Exercises
 - Quadriceps sets, heel slides, straight leg raises, patellar mobilizations, co-contractions (Weeks 0-2)
 - Add heel raises and terminal knee extensions (Weeks 2-8)
 - Activities in brace for first 6 weeks then without brace
 - No weightbearing with flexion > 90° during Phase I
 - Avoid tibial rotation for first 8 weeks to protect the meniscal allograft

Phase II (Weeks 8-12)

- Weightbearing: As tolerated
- Range of Motion Full active ROM
- Therapeutic Exercises
 - Progress to closed chain extension exercises, begin hamstring strengthening
 - o Lunges 0-90°, Leg press 0-90° (flexion only)
 - Proprioception exercises
 - o Begin use of the stationary bicycle

Phase III (Months 3-6)

- Weightbearing: Full weightbearing with normal gait pattern
- Range of Motion Full/Painless ROM
- Therapeutic Exercises
 - Continue with quad and hamstring strengthening
 - o Focus on single-leg strength

- Begin jogging/running
 Plyometrics and sport-specific drills
 Gradual return to athletic activity as tolerated (6 months post-op)
- o Maintenance program for strength and endurance

Comments:			
Frequency:	times per week Duration:	weeks Date:	